arXiv:1209.4579v1 [hep-ph] 20 Sep 2012

From Dirac neutrino masses to baryonic and dark matter asymmetries

Pei-Hong Gu*
Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

We consider an $SU(3)'_c\times SU(2)'_L\times U(1)'_Y$ dark sector, parallel to the $SU(3)_c\times SU(2)_L\times U(1)_Y$ ordinary sector. The hypercharges, baryon numbers and lepton numbers in the dark sector are opposite to those in the ordinary sector. We further introduce three types of messenger sectors: (i) two or more gauge-singlet Dirac fermions, (ii) two or more $[SU(2)_L\times SU(2)'_L]$-bidoublet Higgs scalars, (iii) at least one gauge-singlet Dirac fermion and at least one $[SU(2)_L\times SU(2)'_L]$-bidoublet Higgs scalar. The lepton number conserving decays of the heavy fermion singlet(s) and/or Higgs bidoublet(s) can simultaneously generate a lepton asymmetry in the $[SU(2)_L]$-doublet leptons and an opposite lepton asymmetry in the $[SU(2)'_L]$-doublet leptons to account for the cosmological baryon asymmetry and dark matter relic density, respectively. The lightest dark nucleon as the dark matter particle should have a mass about 5 GeV. By integrating out the heavy fermion singlet(s) and/or Higgs bidoublet(s), we can obtain three light Dirac neutrinos composed of the ordinary and dark neutrinos. If a mirror symmetry is further imposed, our models will not require more unknown parameters than the traditional type-I, type-II or type-I+II seesaw models.

PACS numbers: 14.60.Pq, 98.80.Cq, 95.35.+d, 12.60.Cn, 12.60.Fr

I. INTRODUCTION

Various neutrino oscillation experiments have established the phenomenon of massive and mixing neutrinos. To naturally understand the smallness of neutrino masses, we can consider the seesaw [1] extension of the $SU(3)_c\times SU(2)_L\times U(1)_Y$ standard model (SM). In the conventional seesaw scenario [1–3], the neutrinos have a Majorana nature which, however, has not been experimentally verified. Alternatively, we can build some Dirac seesaw models [4–6] to give the light Dirac neutrinos. Usually, the Dirac seesaw needs more parameters than the Majorana seesaw. In the Majorana or Dirac seesaw context, we can obtain the cosmological baryon asymmetry through the leptogenesis [7–14] or neutrinogenesis [5, 6, 15, 16] mechanism. On the other hand, the dark matter relic density also indicates the necessity of supplementing the SM. It is intriguing that the baryonic and dark matter contribute comparable energy densities to the present universe [17] although they have very different properties. This coincidence can be elegantly explained if the dark matter relic density is an asymmetry between the dark matter and antimatter and has a common origin with the baryon asymmetry. There have been a lot of works studying the asymmetric dark matter scenario [18–45].

In this paper we shall propose a common genesis of the Dirac neutrino masses, the baryon asymmetry and the dark matter relic density. Specifically, we consider an $SU(3)'_c\times SU(2)'_L\times U(1)'_Y$ dark sector, parallel to the ordinary SM sector. The hypercharges, baryon numbers and lepton numbers in the dark sector are opposite to those in the ordinary sector. Besides the ordinary and dark sectors, there are three types of messenger sectors: (i) two or more gauge-singlet Dirac fermions, (ii) two or more $[SU(2)_L\times SU(2)'_L]$-bidoublet Higgs scalars, (iii) at least one gauge-singlet Dirac fermion and at least one $[SU(2)_L\times SU(2)'_L]$-bidoublet Higgs scalar. Through the lepton number conserving decays of the heavy fermion singlet(s) and/or Higgs bidoublet(s), we can simultaneously realize a lepton asymmetry in the $[SU(2)_L]$-doublet leptons and an opposite lepton asymmetry in the $[SU(2)'_L]$-doublet leptons to account for the baryon asymmetry and the dark matter relic density, respectively. The lightest dark nucleon with a determined mass about 5 GeV can serve as the dark matter particle. The dark photon will become massive although the ordinary photon keeps massless. The kinetic mixing between the ordinary and dark photons can result in a testable scattering of the dark nucleons on the ordinary nucleons. Furthermore, we can get a tiny mass term between the ordinary and dark left-handed neutrinos by integrating out the heavy fermion singlet(s) and/or Higgs bidoublet(s). So, the ordinary and dark neutrinos can naturally form the light Dirac neutrinos. Finally, we can impose a mirror symmetry to reduce the parameters. In this case, our models will not contain additional parameters compared with the traditional Majorana seesaw models.

II. THE ORDINARY AND DARK SECTORS

We denote the ordinary quarks, leptons and scalars by

$$q_L(\mathbf{3},\mathbf{2},+\tfrac{1}{3})=\begin{bmatrix}u_L\\ d_L\end{bmatrix},\ d_R(\mathbf{3},\mathbf{1},-\tfrac{2}{3}),\ u_R(\mathbf{3},\mathbf{1},+\tfrac{4}{3}),$$

$$l_L(\mathbf{1},\mathbf{2},-1)=\begin{bmatrix}\nu_L\\ e_L\end{bmatrix},\ e_R(\mathbf{1},\mathbf{1},-2),$$

$$\phi(\mathbf{1},\mathbf{2},-1)=\begin{bmatrix}\phi^0\\ \phi^-\end{bmatrix},\ \delta(\mathbf{1},\mathbf{1},-4),\qquad(1)$$

---

*Electronic address: peihong.gu@mpi-hd.mpg.de

where the first and second numbers in parentheses are the dimensions of the $SU(3)_c$ and $SU(2)_L$ representations, while the third ones are the $U(1)_Y$ hypercharges Y. Accordingly, we define the dark quarks, leptons and scalars as below,

$$\begin{aligned}
&q'_L(\mathbf{3},\mathbf{2},-\tfrac{1}{3})=\left[\begin{array}{c} d'_L \\ u'_L \end{array}\right],\ d'_R(\mathbf{3},\mathbf{1},+\tfrac{2}{3}),\ u'_R(\mathbf{3},\mathbf{1},-\tfrac{4}{3}),\\
&l'_L(\mathbf{1},\mathbf{2},+1)=\left[\begin{array}{c} e'_L \\ \nu'_L \end{array}\right],\ e'_R(\mathbf{1},\mathbf{1},+2),\\
&\phi'(\mathbf{1},\mathbf{2},+1)=\left[\begin{array}{c} \phi'^{+} \\ \phi'^{0} \end{array}\right],\ \delta'(\mathbf{1},\mathbf{1},+4), \qquad (2)
\end{aligned}$$

where the first and second numbers in parentheses are the dimensions of the $SU(3)'_c$ and $SU(2)'_L$ representations, while the third ones are the $U(1)'_Y$ hypercharges Y'. Like the hypercharges, the baryon or lepton numbers of the dark fermions are assumed opposite to those of the ordinary fermions.

We then write down the Lagrangian of the ordinary and dark sectors,

$$\mathcal{L}^{\mathrm{OD}}=\mathcal{L}_K^{\mathrm{OD}}+\mathcal{L}_Y^{\mathrm{OD}}-V^{\mathrm{OD}}\,. \qquad (3)$$

Here the index "OD" is the abbreviation of "ordinary-dark". The kinetic terms include

$$\begin{aligned}
\mathcal{L}_K^{\mathrm{OD}} &= (D_\mu\phi)^\dagger(D^\mu\phi)+(D_\mu\delta)^\dagger(D^\mu\delta)+i\bar{q}_{L_i}\not{D}q_{L_i}\\
&\quad +i\bar{d}_{R_i}\not{D}d_{R_i}+i\bar{u}_{R_i}\not{D}u_{R_i}+i\bar{l}_{L_i}\not{D}l_{L_i}\\
&\quad +i\bar{e}_{R_i}\not{D}e_{R_i}+(D_\mu\phi')^\dagger(D^\mu\phi')+(D_\mu\delta')^\dagger(D^\mu\delta')\\
&\quad +i\bar{q}'_{L_i}\not{D}q'_{L_i}+i\bar{d}'_{R_i}\not{D}d'_{R_i}+i\bar{u}'_{R_i}\not{D}u'_{R_i}+i\bar{l}'_{L_i}\not{D}l'_{L_i}\\
&\quad +i\bar{e}'_{R_i}\not{D}e'_{R_i}-\frac{1}{4}G^a_{\mu\nu}G^{a\mu\nu}-\frac{1}{4}W^a_{\mu\nu}W^{a\mu\nu}\\
&\quad -\frac{1}{4}B_{\mu\nu}B^{\mu\nu}-\frac{1}{4}G'^a_{\mu\nu}G'^{a\mu\nu}-\frac{1}{4}W'^a_{\mu\nu}W'^{a\mu\nu}\\
&\quad -\frac{1}{4}B'_{\mu\nu}B'^{\mu\nu}-\frac{\epsilon}{2}B_{\mu\nu}B'^{\mu\nu}\,, \qquad (4)
\end{aligned}$$

where the covariant derivatives are given by

$$\begin{aligned}
D_\mu &= \partial_\mu-ig_1\frac{Y}{2}B_\mu-ig_2\frac{\tau_a}{2}W^a_\mu\delta_2-ig_3\frac{\lambda_a}{2}G^a_\mu\delta_3\\
&\quad -ig'_1\frac{Y'}{2}B'_\mu-ig'_2\frac{\tau_a}{2}W'^a_\mu\delta'_2-ig'_3\frac{\lambda_a}{2}G'^a_\mu\delta'_3 \ \text{ with}\\
&\quad \delta_2=\left\{\begin{array}{l} 1\ \text{ for } SU(2)_L \text{ doublets}\,,\\ 0\ \text{ for } SU(2)_L \text{ singlets}\,; \end{array}\right.\\
&\quad \delta_3=\left\{\begin{array}{l} 1\ \text{ for } SU(3)_c \text{ triplets}\,,\\ 0\ \text{ for } SU(3)_c \text{ singlets}\,; \end{array}\right. \qquad (5)\\
&\quad \delta'_2=\left\{\begin{array}{l} 1\ \text{ for } SU(2)'_L \text{ doublets}\,,\\ 0\ \text{ for } SU(2)'_L \text{ singlets}\,; \end{array}\right.\\
&\quad \delta'_3=\left\{\begin{array}{l} 1\ \text{ for } SU(3)'_c \text{ triplets}\,,\\ 0\ \text{ for } SU(3)'_c \text{ singlets}\,. \end{array}\right.
\end{aligned}$$

We also read the Yukawa interactions:

$$\begin{aligned}
\mathcal{L}_Y^{\mathrm{OD}} &= -(y_d)_{ij}\bar{q}_{L_i}\tilde{\phi}d_{R_j}-(y_u)_{ij}\bar{q}_{L_i}\phi u_{R_j}\\
&\quad -(y_e)_{ij}\bar{l}_{L_i}\tilde{\phi}e_{R_j}-y_\delta\delta\bar{e}_Re^c_R-(y'_d)_{ij}\bar{q}'_{L_i}\tilde{\phi}'d'_{R_j}\\
&\quad -(y'_u)_{ij}\bar{q}'_{L_i}\phi'u'_{R_j}-(y'_e)_{ij}\bar{l}'_{L_i}\tilde{\phi}'e'_{R_j}\\
&\quad -y_{\delta'}\delta'\bar{e}'_Re'^c_R+\mathrm{H.c.}\,, \qquad (6)
\end{aligned}$$

and the scalar potential:

$$\begin{aligned}
V^{\mathrm{OD}} &= \mu^2_\phi\phi^\dagger\phi+\lambda_\phi(\phi^\dagger\phi)^2+\mu^2_\delta\delta^\dagger\delta+\lambda_\delta(\delta^\dagger\delta)^2\\
&\quad +\lambda_{\phi\delta}\phi^\dagger\phi\delta^\dagger\delta+\mu^2_{\phi'}\phi'^\dagger\phi'+\lambda_{\phi'}(\phi'^\dagger\phi')^2\\
&\quad +\mu^2_{\delta'}\delta'^\dagger\delta'+\lambda_{\delta'}(\delta'^\dagger\delta')^2+\lambda_{\phi'\delta'}\phi'^\dagger\phi'\delta'^\dagger\delta'\\
&\quad +2\lambda_{\phi\phi'}\phi^\dagger\phi\phi'^\dagger\phi'+2\lambda_{\phi\delta'}\phi^\dagger\phi\delta'^\dagger\delta'\\
&\quad +2\lambda_{\delta\phi'}\delta^\dagger\delta\phi'^\dagger\phi'+2\lambda_{\delta\delta'}\delta^\dagger\delta\delta'^\dagger\delta'\,. \qquad (7)
\end{aligned}$$

The symmetry breaking pattern is expected to be

$$\begin{aligned}
SU(3)_c\times SU(2)_L\times U(1)_Y &\longrightarrow SU(3)_c\times U(1)_{em}\,,\\
SU(3)'_c\times SU(2)'_L\times U(1)'_Y &\longrightarrow SU(3)'_c\times U(1)'_{em}\\
&\longrightarrow SU(3)'_c\,. \qquad (8)
\end{aligned}$$

For this purpose, the ordinary and dark scalars should develop their vacuum expectation values (VEVs) as below,

$$\begin{aligned}
\langle\phi\rangle &= \left[\begin{array}{c} \langle\phi^0\rangle \\ 0 \end{array}\right],\ \ \langle\delta\rangle=0\,,\\
\langle\phi'\rangle &= \left[\begin{array}{c} 0 \\ \langle\phi'^0\rangle \end{array}\right],\ \ \langle\delta'\rangle\neq 0\,. \qquad (9)
\end{aligned}$$

This means the ordinary photon γ will keep massless while the mirror photon γ' will become massive.

It is straightforward to read the fermion masses in the ordinary sector,

$$\begin{aligned}
\mathcal{L} &\supset -m_d\bar{d}_Ld_R-m_u\bar{u}_Lu_R-m_e\bar{e}_Le_R+\mathrm{H.c.}\ \text{ with}\\
&\quad m_d=y_d\langle\phi\rangle\,,\ m_u=y_u\langle\phi\rangle\,,\ m_e=y_e\langle\phi\rangle\,, \qquad (10)
\end{aligned}$$

and the fermion masses in the dark sector,

$$\begin{aligned}
\mathcal{L} &\supset -m_{d'}\bar{d}'_Rd'_L-m_{u'}\bar{u}'_Ru'_L-m_{e'}\bar{e}'_Re'_L-\delta m_{e'}\bar{e}'_Re'^c_R\\
&\quad +\mathrm{H.c.}\ \text{ with }\ m_{d'}=y_{d'}\langle\phi'\rangle\,,\ m_{u'}=y_{u'}\langle\phi'\rangle\,,\\
&\quad m_{e'}=y_{e'}\langle\phi'\rangle\,,\ \delta m_{e'}=y_{\delta'}\langle\delta'\rangle\,. \qquad (11)
\end{aligned}$$

The dark charged leptons should be the so-called quasi-Dirac fermions for $\delta m_{e'}\ll m_{e'}$. In the ordinary sector, the quark masses m_u and m_d are much smaller than the hadronic scale Λ_{QCD} so that they can only have a negligible contribution to the nucleon masses,

$$m_p\simeq m_n\simeq 1\,\mathrm{GeV}=m_N\,. \qquad (12)$$

In the dark sector, the quark masses $m_{u'}$ and $m_{d'}$ may be sufficiently larger than the hadronic scale $\Lambda_{\mathrm{QCD}'}$. In this



FIG. 1: The type-I Dirac seesaw for generating the masses between the ordinary left-handed neutrinos ν_L and the dark left-handed neutrinos ν'_L.

case, the dark nucleon masses can approximately equal to the sum of the dark quark masses,

$$m_{p'} = 2m_{u'} + m_{d'}\,, \quad m_{n'} = 2m_{d'} + m_{u'}\,. \tag{13}$$

As we will show in the following, our completed models also contain a messenger sector. We will refer to $\mathcal{L}^{\mathrm{M}}$ (with the index "M" being the abbreviation of "messenger") as the Lagrangian involving the messenger fields. Note that our models will not have any baryon or lepton number violating interactions except the $SU(2)_L$ and $SU(2)'_L$ sphaleron processes [46, 47].

III. THE MODEL WITH GAUGE-SINGLET DIRAC FERMIONS

In this sector, we will give the completed model with two or more gauge-singlet Dirac fermions. The Lagrangian involving the fermion singlets (FSs) should be

$$\mathcal{L}^{\mathrm{M}} \equiv \mathcal{L}^{\mathrm{FS}} = \mathcal{L}^{\mathrm{FS}}_K + \mathcal{L}^{\mathrm{FS}}_Y + \mathcal{L}^{\mathrm{FS}}_m\,, \tag{14}$$

with the kinetic terms:

$$\mathcal{L}^{\mathrm{FS}}_K = i\bar{N}_{R_i}\not{\partial}N_{R_i} + i\bar{N}'_{R_i}\not{\partial}N'_{R_i}\,, \tag{15}$$

the Yukawa terms:

$$\mathcal{L}^{\mathrm{FS}}_Y = -(y_N)_{ij}\bar{l}_{L_i}\phi N_{R_j} - (y_{N'})_{ij}\bar{l}'_{L_i}\phi' N'_{R_j} + \mathrm{H.c.}\,, \tag{16}$$

and the mass terms:

$$\mathcal{L}^{\mathrm{FS}}_m = -(M_N)_{ij}\bar{N}'^c_{R_i}N_{R_j} + \mathrm{H.c.}\,. \tag{17}$$

Here we have introduced two types of gauge-singlet right-handed fermions:

$$\left\{\begin{array}{lll} N_R(\mathbf{1},\mathbf{1},0)(\mathbf{1},\mathbf{1},0) & \text{with a lepton number} & +1\,,\\ N'_R(\mathbf{1},\mathbf{1},0)(\mathbf{1},\mathbf{1},0) & \text{with a lepton number} & -1\,,\end{array}\right. \tag{18}$$

where the first and second parentheses being the quantum numbers under the ordinary and dark gauge groups, respectively. Note that other gauge-invariant Yukawa and mass terms have been forbidden as a result of the lepton number conservation. After choosing a proper base, the fermion singlets can have a diagonal and real mass matrix:

$$M_N = \mathrm{diag}\{M_{N_1}\,,\ M_{N_2}\,,...\}\,. \tag{19}$$

We then can define the Dirac fermions:

$$N_i = N_{R_i} + N'^c_{R_i}\,. \tag{20}$$

A. Dirac neutrino masses

After the $[SU(2)_L]$-doublet Higgs scalar ϕ and the $[SU(2)'_L]$-doublet Higgs scalar ϕ' develop their VEVs, the ordinary and dark left-handed neutrinos as well as the gauge-singlet right-handed fermions will have a mass matrix as below,

$$\mathcal{L} \supset -\left[\begin{array}{cc}\bar{\nu}_L & \bar{N}'^c_R\end{array}\right]\left[\begin{array}{cc}0 & y_N\langle\phi\rangle\\ y^T_{N'}\langle\phi'\rangle & M_N\end{array}\right]\left[\begin{array}{c}\nu'^c_L\\ N_R\end{array}\right] + \mathrm{H.c.} \tag{21}$$

We can block diagonalize the above mass matrix to be

$$\mathcal{L} \supset -\left[\begin{array}{cc}\bar{\nu}_L & \bar{N}'^c_R\end{array}\right]\left[\begin{array}{cc}m_\nu & 0\\ 0 & M_N\end{array}\right]\left[\begin{array}{c}\nu'^c_L\\ N_R\end{array}\right] + \mathrm{H.c.}\ \ \text{with}$$

$$m_\nu = -y_N\frac{\langle\phi\rangle\langle\phi'\rangle}{M_N}y^T_{N'} \equiv m^{\mathrm{I}}_\nu\,, \tag{22}$$

if the seesaw condition is satisfied, i.e.

$$y_N\langle\phi\rangle\,,\ y_{N'}\langle\phi'\rangle \ll M_N\,. \tag{23}$$

This means the ordinary and dark left-handed neutrinos will form the light Dirac neutrinos $\nu = \nu_L + \nu'^c_L$, while the gauge-singlet right-handed fermions will form the heavy Dirac fermions $N = N_R + N'^c_R$. Note that the Dirac neutrino mass matrix (22) will have one nonzero eigenvalue if there is only one gauge-singlet Dirac fermion. We thus need two or more gauge-singlet Dirac fermions to explain the neutrino oscillation data. Analogous to the usual type-I seesaw [1] formula of the Majorana neutrino masses, we shall refer to the formula (22) of the Dirac neutrino masses to be the type-I Dirac seesaw. The relevant diagram is shown in Fig. 1.

B. Ordinary and dark lepton asymmetries

As shown in Fig. 2, the lepton number conserving decays of the gauge-singlet Dirac fermions N_i can simultaneously generate a lepton asymmetry in the $[SU(2)_L]$-doublet leptons l_L and an opposite lepton asymmetry in the $[SU(2)'_L]$-doublet leptons l'_L, i.e.

$$\eta_{l_L} = -\eta_{l'_L} \propto \varepsilon_{N_i}\,. \tag{24}$$



FIG. 2: The lepton number conserving decays of the heavy gauge-singlet Dirac fermions $N_i = N_{R_i} + N'^c_{R_i}$ in the type-I Dirac seesaw scenario. We need at leat two fermion singlets to generate a nonzero lepton asymmetry in the $[SU(2)_L]$-doublet leptons l_L and an opposite lepton asymmetry in the $[SU(2)'_L]$-doublet leptons l'_L. The CP conjugation is not shown for simplicity.

Here the CP asymmetry ε_{N_i} can be calculated by

$$\begin{aligned}\varepsilon_{N_i} &= \frac{\Gamma_{N_i\to l_L\phi^*} - \Gamma_{N_i^c\to l_L^c\phi}}{\Gamma_{N_i}} = \frac{\Gamma_{N_i^c\to l'_L\phi'^*} - \Gamma_{N_i\to l'^c_L\phi'}}{\Gamma_{N_i}}\\ &= -\frac{1}{4\pi}\sum_{j\neq i}\frac{\mathrm{Im}\left[(y_N^\dagger y_N)_{ij}(y_{N'}^\dagger y_{N'})_{ij}\right]}{(y_N^\dagger y_N)_{ii} + (y_{N'}^\dagger y_{N'})_{ii}}\frac{M_{N_i}M_{N_j}}{M_{N_j}^2 - M_{N_i}^2}\,,\end{aligned} \tag{25}$$

with Γ_{N_i} being the decay width:

$$\begin{aligned}\Gamma_{N_i} &= \Gamma_{N_i\to l_L\phi^*} + \Gamma_{N_i\to l'^c_L\phi'} = \Gamma_{N_i^c\to l_L^c\phi} + \Gamma_{N_i^c\to l'_L\phi'^*}\\ &= \frac{1}{16\pi}[(y_N^\dagger y_N)_{ii} + (y_{N'}^\dagger y_{N'})_{ii}]M_{N_i}\,.\end{aligned} \tag{26}$$

We should keep in mind that at least two gauge-singlet Dirac fermions are necessary to induce a nonzero CP asymmetry.

If the gauge-singlet Dirac fermions have a hierarchical mass spectrum, i.e $M^2_{N_i} \ll M^2_{N_j}$, we can simplify the CP asymmetry (25) to be

$$\varepsilon_{N_i} \simeq \frac{1}{4\pi}\frac{\mathrm{Im}\left[(y_N^\dagger m_\nu y_{N'}^*)_{ii}\right]}{(y_N^\dagger y_N)_{ii} + (y_{N'}^\dagger y_{N'})_{ii}}\frac{M_{N_i}}{\langle\phi\rangle\langle\phi'\rangle}\,. \tag{27}$$

Similar to the Davidson-Ibarra bound [10] in the type-I Majorana seesaw scenario, the above CP asymmetry should have an upper bound:

$$\begin{aligned}|\varepsilon_{N_i}| &\leq \frac{1}{8\pi}\frac{\left|\mathrm{Im}\left[(y_N^\dagger m_\nu y_{N'}^*)_{ii}\right]\right|}{\sqrt{(y_N^\dagger y_N)_{ii}(y_{N'}^\dagger y_{N'})_{ii}}}\frac{M_{N_i}}{\langle\phi\rangle\langle\phi'\rangle}\\ &< \frac{1}{8\pi}\frac{M_{N_i}m_\nu^{\max}}{\langle\phi\rangle\langle\phi'\rangle}\,.\end{aligned} \tag{28}$$

Here $m_\nu^{\max}$ is the maximal eigenvalue of the neutrino mass matrix m_ν. Alternatively, the CP asymmetry (25) can be resonantly enhanced [9] if the gauge-singlet Dirac fermions have a quasi-degenerate mass spectrum, i.e $M^2_{N_i} \simeq M^2_{N_j} \gg |M^2_{N_i} - M^2_{N_j}|$.

IV. THE MODEL WITH $[SU(2)_L \times SU(2)'_L]$-BIDOUBLET HIGGS SCALARS

In this sector, we will give the completed model with two or more $[SU(2)_L \times SU(2)'_L]$-bidoublet Higgs scalars:

$$\Sigma_a(\mathbf{1},\mathbf{2},-\frac{1}{2})(\mathbf{1},\mathbf{2},+\frac{1}{2}) = \left[\begin{array}{cc}\bar{\sigma}^+ & \sigma^0\\ \bar{\sigma}^0 & \sigma^-\end{array}\right]_a\,. \tag{29}$$

Here the first and second parentheses stand for the quantum numbers under the ordinary and dark gauge groups, respectively. The Lagrangian involving the Higgs bidoublets (HBs) should be

$$\mathcal{L}^{\mathrm{M}} \equiv \mathcal{L}^{\mathrm{HB}} = \mathcal{L}_K^{\mathrm{HB}} + \mathcal{L}_Y^{\mathrm{HB}} - V^{\mathrm{HB}}\,, \tag{30}$$

where the kinetic terms include

$$\begin{aligned}\mathcal{L}_K^{\mathrm{HB}} &= \mathrm{Tr}[(D_\mu\Sigma_a)^\dagger(D_\mu\Sigma_a)] \ \text{ with}\\ &\qquad D_\mu\Sigma_a = \partial_\mu\Sigma_a + i\frac{1}{2}g_1B_\mu\Sigma_a - ig_2\frac{\tau_a}{2}W^a_\mu\Sigma_a\\ &\qquad\qquad -i\frac{1}{2}g'_1B'_\mu\Sigma_a - ig'_2\Sigma_a\frac{\tau_a^T}{2}W'^a_\mu\,,\end{aligned} \tag{31}$$

the Yukawa couplings contain

$$\mathcal{L}_Y^{\mathrm{HB}} = -f_a\bar{l}_L\Sigma_a l'^c_L + \mathrm{H.c.}\,, \tag{32}$$



FIG. 3: The type-II Dirac seesaw for generating the masses between the ordinary left-handed neutrinos ν_L and the dark left-handed neutrinos ν'_L.

and the scalar potential is

$$\begin{aligned} V^{\rm HB} &= (M_\Sigma^2)_{ab}{\rm Tr}(\Sigma_a^\dagger\Sigma_b) + (\lambda_\Sigma)_{abcd}{\rm Tr}(\Sigma_a^\dagger\Sigma_b){\rm Tr}(\Sigma_c^\dagger\Sigma_d)\\ &\quad +[(\lambda_{\phi\Sigma})_{ab}\phi^\dagger\phi + (\lambda_{\phi'\Sigma})_{ab}\phi'^\dagger\phi']{\rm Tr}(\Sigma_a^\dagger\Sigma_b)\\ &\quad +[(\lambda_{\delta\Sigma})_{ab}\delta^\dagger\delta + (\lambda_{\delta'\Sigma})_{ab}\delta'^\dagger\delta']{\rm Tr}(\Sigma_a^\dagger\Sigma_b)\\ &\quad +\rho_a\phi^\dagger\Sigma_a\phi'^* + {\rm H.c.}\,. \end{aligned} \tag{33}$$

Without loss of generality, we can choose a base to take

$$M_\Sigma^2 = {\rm diag}\{M_{\Sigma_1}^2\,,\ M_{\Sigma_2}^2\,,\ ...\}\,,\ \rho_1=\rho_1^*\,,\ \rho_2=\rho_2^*\,, ... \tag{34}$$

Note that the above Yukawa couplings and scalar potential will exactly conserve the lepton number because the Higgs bidoublets Σ_a don't carry any lepton numbers.

A. Dirac neutrino masses

After the $[SU(2)_L]$-doublet Higgs scalar ϕ and the $[SU(2)'_L]$-doublet Higgs scalar ϕ' acquire their VEVs, the heavy $[SU(2)_L\times SU(2)'_L]$-bidoublet Higgs scalars Σ_a can pick up the seesaw-suppressed VEVs:

$$\langle\Sigma_a\rangle = \left[\begin{array}{cc} 0 & \frac{1}{\sqrt{2}}u_a \\ 0 & 0 \end{array}\right] \quad \text{with} \quad u_a \simeq -\frac{\rho_a v v'}{\sqrt{2}M_{\Sigma_a}^2} \ll v, v'\,. \tag{35}$$

We hence can naturally obtain the light Dirac neutrinos composed of the ordinary left-handed neutrinos ν_L and the dark left-handed neutrinos ν'_L, i.e.

$$\begin{aligned} \mathcal{L} &\supset -m_\nu\bar{\nu}_L\nu'^c_L + {\rm H.c.} \quad \text{with}\\ & \quad m_\nu = \sum_a f_a\langle\Sigma_a\rangle \equiv \sum_a m_\nu^{{\rm II}a} \equiv m_\nu^{\rm II}\,. \end{aligned} \tag{36}$$

The above mechanism for the Dirac neutrino masses is very similar to the usual type-II seesaw [2] for the Majorana neutrino masses. So, it may be named as the type-II Dirac seesaw. We show the relevant diagram in Fig. 3.

B. Ordinary and dark lepton asymmetries

From Fig. 4, it is straightforward to see that the lepton number conserving decays of the $[SU(2)_L\times SU(2)'_L]$-bidoublet Higgs scalars Σ_a can generate a lepton asymmetry in the $[SU(2)_L]$-doublet leptons l_L and an opposite lepton asymmetry in the $[SU(2)'_L]$-doublet leptons l'_L, i.e.

$$\eta_{l_L} = -\eta_{l'_L} \propto \varepsilon_{\Sigma_a}\,, \tag{37}$$

where ε_a is the CP asymmetry:

$$\begin{aligned} \varepsilon_{\Sigma_a} &= \frac{\Gamma_{\Sigma_a\to l_L l'_L} - \Gamma_{\Sigma_a^*\to l_L^c l'^c_L}}{\Gamma_{\Sigma_a}}\\ &= \frac{\Gamma_{\Sigma_a^*\to\phi^*\phi'^*} - \Gamma_{\Sigma_a\to\phi\phi'}}{\Gamma_{\Sigma_a}}\,, \end{aligned} \tag{38}$$

with Γ_{Σ_a} being the decay width:

$$\begin{aligned} \Gamma_{\Sigma_a} &= \Gamma_{\Sigma_a\to l_L l'_L} + \Gamma_{\Sigma_a\to\phi\phi'}\\ &= \Gamma_{\Sigma_a^*\to l_L^c l'^c_L} + \Gamma_{\Sigma_a^*\to\phi^*\phi'^*}\,. \end{aligned} \tag{39}$$

We can calculate the decay width at tree level,

$$\Gamma_{\Sigma_a} = \frac{1}{16\pi}\left[{\rm Tr}\left(f_a^\dagger f_a\right) + \frac{\rho_a^2}{M_{\Sigma_a}^2}\right]M_{\Sigma_a}\,, \tag{40}$$

and the CP asymmetry at one-loop order,

$$\varepsilon_{\Sigma_a} = -\frac{1}{4\pi}\sum_{b\neq a}\frac{{\rm Im}\left[{\rm Tr}\left(f_b^\dagger f_a\right)\right]}{{\rm Tr}\left(f_a^\dagger f_a\right) + \frac{\rho_a^2}{M_{\Sigma_a}^2}}\frac{\rho_b\rho_a}{M_{\Sigma_b}^2 - M_{\Sigma_a}^2}\,. \tag{41}$$

Note that at least two $[SU(2)_L\times SU(2)'_L]$-bidoublet Higgs scalars should be introduced to generate a nonzero CP asymmetry.

Obviously, the above CP asymmetry can be resonantly enhanced if the $[SU(2)_L\times SU(2)'_L]$-bidoublet Higgs scalars have a quasi-degenerate mass spectrum, i.e $M_{\Sigma_a}^2 \simeq M_{\Sigma_b}^2 \gg |M_{\Sigma_a}^2 - M_{\Sigma_b}^2|$. In the hierarchical case with $M_{\Sigma_a}^2 \ll M_{\Sigma_b}^2$, the CP asymmetry (41) can be simplified by

$$\varepsilon_{\Sigma_a} \simeq -\frac{1}{4\pi}\frac{\sum_{b\neq a}{\rm Im}\left[{\rm Tr}\left(m_\nu^{{\rm II}b\dagger}m_\nu^{{\rm II}a}\right)\right]}{{\rm Tr}\left(f_a^\dagger f_a\right) + \frac{\rho_a^2}{M_{\Sigma_a}^2}}\frac{M_{\Sigma_a}^2}{\langle\phi\rangle^2\langle\phi'\rangle^2}\,, \tag{42}$$

and then have an upper bound:

$$\begin{aligned} |\varepsilon_{\Sigma_a}| &\leq \frac{1}{8\pi}\frac{\left|\sum_{b\neq a}{\rm Im}\left[{\rm Tr}\left(m_\nu^{{\rm II}b\dagger}m_\nu^{{\rm II}a}\right)\right]\right|}{\sqrt{{\rm Tr}\left(f_a^\dagger f_a\right)\frac{\rho_a^2}{M_{\Sigma_a}^2}}}\frac{M_{\Sigma_a}^2}{\langle\phi\rangle^2\langle\phi'\rangle^2}\\ &= \frac{1}{8\pi}\frac{\left|\sum_{b\neq a}{\rm Im}\left[{\rm Tr}\left(m_\nu^{{\rm II}b\dagger}m_\nu^{{\rm II}a}\right)\right]\right|}{\sqrt{{\rm Tr}\left(m_\nu^{{\rm II}a\dagger}m_\nu^{{\rm II}a}\right)}}\frac{M_{\Sigma_a}}{\langle\phi\rangle\langle\phi'\rangle}\\ &< \frac{1}{8\pi}\frac{M_{\Sigma_a}m_\nu^{{\rm II}b{\rm max}}}{\langle\phi\rangle\langle\phi'\rangle}\,. \end{aligned} \tag{43}$$



FIG. 4: The lepton number conserving decays of the heavy $[SU(2)_L \times SU(2)'_L]$-bidoublet Higgs scalars Σ_a in the type-II Dirac seesaw scenario. We need at leat two Higgs bidoublets to generate a nonzero lepton asymmetry in the $[SU(2)_L]$-doublet leptons l_L and an opposite lepton asymmetry in the $[SU(2)'_L]$-doublet leptons l'_L. The CP conjugation is not shown for simplicity.

Here $m_\nu^{\mathrm{II}b\max}$ is the maximal eigenvalue of the mass matrix $\sum_{b\neq a} m_\nu^{\mathrm{II}b}$. Unless there is a large cancelation between the mass matrix $m_\nu^{\mathrm{II}a}$ and the mass matrix $\sum_{b\neq a} m_\nu^{\mathrm{II}b}$, the eigenvalue $m_\nu^{\mathrm{II}b\max}$ will not be much bigger than the largest neutrino mass $m_\nu^{\max}$. So, we can roughly constrain

$$|\varepsilon_{\Sigma_a}| \;<\; \frac{1}{8\pi}\frac{M_{\Sigma_a} m_\nu^{\max}}{\langle\phi\rangle\langle\phi'\rangle}\,. \tag{44}$$

V. THE MODEL WITH GAUGE-SINGLET DIRAC FERMION(S) AND $[SU(2)_L \times SU(2)'_L]$-BIDOUBLET HIGGS SCALAR(S)

In this sector, we will give the completed model with at least one gauge-singlet Dirac fermion and at least one $[SU(2)_L \times SU(2)'_L]$-bidoublet Higgs scalar. By taking the notations in the previous sections, the Lagrangian involving the fermion singlet(s) and the Higgs bidoublet(s) can be described by

$$\mathcal{L}^{\mathrm{M}} \equiv \mathcal{L}^{\mathrm{FS}} + \mathcal{L}^{\mathrm{HB}}\,. \tag{45}$$

A. Dirac neutrino masses

The ordinary and dark left-handed neutrinos as well as the gauge-singlet right-handed fermions should have the following mass matrix:

$$\mathcal{L} \supset -\left[\,\bar{\nu}_L \;\; \bar{N}'^c_R\,\right]\left[\begin{array}{cc}\sum_a f_a\langle\Sigma_a\rangle & y_N\langle\phi\rangle\\ y^T_{N'}\langle\phi'\rangle & M_N\end{array}\right]\left[\begin{array}{c}\nu'^c_L\\ N_R\end{array}\right] \\ +\mathrm{H.c.}\,. \tag{46}$$

Obviously, the ordinary left-handed neutrinos ν_L and the dark left-handed neutrinos ν'_L will form three light Dirac neutrinos as their mass term is just a sum of the type-I Dirac seesaw (22) and the type-II Dirac seesaw (36), i.e.

$$m_\nu = m_\nu^{\mathrm{I}} + m_\nu^{\mathrm{II}}\,. \tag{47}$$

B. Ordinary and dark lepton asymmetries

As shown in Figs. 5 and 6, the lepton number conserving decays of the heavy gauge-singlet Dirac fermions N_i and/or $[SU(2)_L \times SU(2)'_L]$-bidoublet Higgs scalars Σ_a can simultaneously generate a lepton asymmetry in the $[SU(2)_L]$-doublet leptons l_L and an opposite lepton asymmetry in the $[SU(2)'_L]$-doublet leptons l'_L. The relevant CP asymmetries should be

$$\begin{aligned}\varepsilon_{N_i} \;=\; & -\frac{1}{4\pi}\frac{1}{(y^\dagger_N y_N)_{ii} + (y^\dagger_{N'} y_{N'})_{ii}}\\ & \times\Bigg\{\sum_{j\neq i}\mathrm{Im}\left[(y^\dagger_N y_N)_{ij}(y^\dagger_{N'} y_{N'})_{ij}\right]\frac{M_{N_i}M_{N_j}}{M^2_{N_j}-M^2_{N_i}}\\ & +2\,\mathrm{Im}\left[(y^\dagger_N f_a y^*_{N'})_{ii}\right]\frac{\rho_a}{M_{N_i}}\\ & \times\left[1-\frac{M^2_{\Sigma_a}}{M^2_{N_i}}\ln\left(1+\frac{M^2_{N_i}}{M^2_{\Sigma}}\right)\right]\Bigg\}\,,\end{aligned} \tag{48}$$



FIG. 5: The lepton number conserving decays of the heavy gauge-singlet Dirac fermions $N_i = N_{R_i} + N'^c_{R_i}$ in the type-I+II Dirac seesaw scenario. If there is only one fermion singlet, the self-energy correction will not contribute to the generation of the lepton asymmetries in the $[SU(2)_L]$-doublet leptons l_L and the $[SU(2)'_L]$-doublet leptons l'_L. The CP conjugation is not shown for simplicity.

and

$$\begin{aligned}\varepsilon_{\Sigma_a} &= -\frac{1}{4\pi}\frac{1}{\mathrm{Tr}(f^\dagger_a f_a)+\frac{\rho^2_a}{M^2_{\Sigma_a}}}\left\{\sum_{b\neq a}\frac{\mathrm{Im}\left[\mathrm{Tr}\left(f^\dagger_b f_a\right)\right]\rho_b\rho_a}{M^2_{\Sigma_b}-M^2_{\Sigma_a}}\right.\\ &\quad\left.+\sum_i \mathrm{Im}\left[(y^\dagger_N f_a y^*_{N'})_{ii}\right]\frac{\rho_a M_{N_i}}{M^2_{\Sigma_a}}\ln\left(1+\frac{M^2_{\Sigma_a}}{M^2_{N_i}}\right)\right\}.\end{aligned} \tag{49}$$

Clearly, the vertex correction should be the unique source for the nonzero CP asymmetry (48) if there is only one gauge-singlet Dirac fermion. As for the CP asymmetry (49), it will not be affected by the self-energy correction if we only introduce one $[SU(2)_L \times SU(2)'_L]$-bidoublet Higgs scalar.

Similar to those in the pure type-I and type-II Dirac seesaw models, the CP asymmetries (48) and (49) can be simplified in the hierarchical cases, i.e.

$$\begin{aligned}\varepsilon_{N_i} &\simeq \frac{1}{4\pi}\frac{\mathrm{Im}\left[(y^\dagger_N m_\nu y^*_{N'})_{ii}\right]}{(y^\dagger_N y_N)_{ii}+(y^\dagger_{N'} y_{N'})_{ii}}\frac{M_{N_i}}{\langle\phi\rangle\langle\phi'\rangle}\\ &\quad \text{for}~~ M^2_{N_i} \ll M^2_{N_j}\,, M^2_{\Sigma_a}\,, \end{aligned}\tag{50}$$

$$\begin{aligned}\varepsilon_{\Sigma_a} &\simeq -\frac{1}{4\pi}\frac{\mathrm{Im}\left\{\mathrm{Tr}[(m^{\mathrm{I}}_\nu + \sum_{b\neq a} m^{\mathrm{II}b}_\nu)^\dagger m^{\mathrm{II}a}_\nu]\right\}}{\mathrm{Tr}(f^\dagger_a f_a)+\frac{\rho^2_a}{M^2_{\Sigma_a}}}\frac{M^2_{\Sigma_a}}{\langle\phi\rangle^2\langle\phi'\rangle^2}\\ &\quad \text{for}~~ M^2_{\Sigma_a} \ll M^2_{\Sigma_b}\,,~ M^2_{N_i}\,, \end{aligned}\tag{51}$$

and have the upper bounds given in Eqs. (28) and (44).

VI. PREDICTION, CONSTRAINT AND IMPLICATION

The $SU(2)_L$ sphaleron processes [47] will partially transfer the lepton asymmetry η_{l_L} to a baryon asymmetry in the ordinary sector,

$$\eta_B = -\frac{28}{79}\eta_{l_L}\,. \tag{52}$$

On the other hand, the lepton asymmetry $\eta_{l'_L}$ will result in a baryon asymmetry in the dark sector,

$$\eta'_B = -\frac{28}{79}\eta_{l'_L}\,, \tag{53}$$

through the $SU(2)'_L$ sphaleron processes [1]. So, the correlation between the baryon asymmetries in the ordinary and dark sectors should be

$$\eta'_B = -\eta_B ~~\text{for}~~ \eta_{l_L} = -\eta_{l'_L}\,. \tag{54}$$

If the dark matter relic density is dominated by the dark proton and/or neutron, we should have

$$\begin{aligned}\Omega_B h^2 : \Omega_{DM} h^2 &= m_p \eta_B : m_{p'(n')}(-\eta'_B) = m_p : m_{p'(n')}\\ &\simeq 1:5\,,\end{aligned} \tag{55}$$

---

[1] After the $U(1)'_{em}$ symmetry is broken, there will be a lepton number violating mass term of the dark right-handed charged leptons, i.e the $\delta m_{e'}$ term in Eq. (11). The $U(1)'_{em}$ symmetry breaking and then the lepton number violation should appear at a lower temperature, where the $SU(2)'_L$ sphaleron processes have become very weak, to avoid the washout of the dark baryon asymmetry.



FIG. 6: The lepton number conserving decays of the heavy $[SU(2)_L \times SU(2)_L']$-bidoublet Higgs scalars Σ_a in the type-I+II Dirac seesaw scenario. If there is only one Higgs bidoublet, the self-energy correction will not contribute to the generation of the lepton asymmetries in the $[SU(2)_L]$-doublet leptons l_L and the $[SU(2)_L']$-doublet leptons l_L'. The CP conjugation is not shown for simplicity.

to fit the cosmological observations. In this case, the dark matter mass should be determined by [2]

$$m_{p'(n')} \simeq 5\,\mathrm{GeV}\,. \tag{56}$$

The Big-Bang Nucleosynthesis (BBN) stringently restricts the existence of the new relativistic degrees of freedom. The constraint on the new degrees of freedom is conventionally quoted as ΔN_ν, the effective number of additional light neutrinos. The seven-year WMAP observation has measured [17]

$$\Delta N_\nu = 1.34^{+0.86}_{-0.88}(68\%\,\mathrm{CL})\,. \tag{57}$$

Assuming the dark charged fermions and the dark photon heavy enough, we need only consider the dark left-handed neutrinos which form the light Dirac neutrinos with the ordinary left-handed neutrinos. We can estimate the decoupling temperature of the dark neutrinos by [49]

$$\begin{aligned}&\left(\frac{g_2'/\cos\theta_W'}{g_2/\cos\theta_W}\right)^4\left(\frac{\langle\phi\rangle}{\langle\phi'\rangle}\right)^4 G_F^2 T^5 = H(T)\\ &\Rightarrow T \simeq 8.6\,\mathrm{GeV}\left(\frac{g_*}{100}\right)^{\frac{1}{6}}\left(\frac{\langle\phi'\rangle}{500\,\langle\phi\rangle}\right)^{\frac{4}{3}}\left(\frac{g_2/\cos\theta_W}{g_2'/\cos\theta_W'}\right)^{\frac{4}{3}}.\end{aligned} \tag{58}$$

Here $G_F = 1.16637\times 10^{-5}\,\mathrm{GeV}^{-2}$ is the Fermi constant [50],

$$\tan\theta_W = \frac{g_1}{g_2}\,, \quad \tan\theta_W' = \frac{g_1'}{g_2'}\,, \tag{59}$$

are the Weinberg angles in the ordinary and dark sectors, while

$$H(T) = \left[\frac{8\pi^3 g_*(T)}{90}\right]^{\frac{1}{2}}\frac{T^2}{M_{\mathrm{Pl}}}\,, \tag{60}$$

is the Hubble constant with $M_{\mathrm{Pl}} \simeq 1.22\times 10^{19}\,\mathrm{GeV}$ being the Planck mass and $g_*(T)$ being the relativistic degrees of freedom. Obviously, the dark neutrinos can decouple before the BBN for a proper choice of the parameters in the dark sector.

The Higgs portal will give a mass mixing between the ordinary Higgs boson h and the dark Higgs boson h', the vacuum expectation value(s) of the $[SU(2)_L \times SU(2)_L']$-bidoublet Higgs scalar(s) will give a mass mixing between the ordinary Z boson and the dark Z' boson, while the kinetic mixing between the $U(1)_Y$ and $U(1)'_Y$ gauge fields will give a kinetic mixing between the the ordinary massless photon γ and the dark massive photon γ'. All of the $h-h'$ mixing, the $Z-Z'$ mixing and the $\gamma-\gamma'$ mixing can mediate a scattering of the dark nucleons on the ordinary nucleons. However, only the scattering induced by the $\gamma-\gamma'$ mixing can arrive at a testable level. The detailed studies can be found in [27].

---

[2] The sphaleron processes in the ordinary and dark sectors could work at different temperatures. For example, the sphalerons in the ordinary sector can keep in equilibrium till a lower temperature T_{sph} after those in the dark sector go out of equilibrium at a higher temperature $T_{\mathrm{sph}'}$. In Eqs. (52) and (53), we have assumed all of the lepton asymmetries to be produced during the epoch that the ordinary and dark sphalerons are both active. Alternatively, the generation of the lepton asymmetries can cross the critical temperature $T_{\mathrm{sph}'}$. In this case, the lepton asymmetries generated after the temperature $T_{\mathrm{sph}'}$ can contribute to the ordinary baryon asymmetry but can not affect the dark baryon asymmetry. This is like the mechanism in the νMSM model for generating a large lepton asymmetry but a small baryon asymmetry [48]. For a proper choice of the masses and couplings of the decaying gauge-singlet Dirac fermion(s) and/or $[SU(2)_L \times SU(2)_L']$-bidoublet Higgs scalar(s), the dark baryon asymmetry can be smaller than the ordinary baryon asymmetry. In consequence, the lightest dark nucleon as the dark matter particle can have a mass bigger than 5 GeV.

VII. MIRROR SYMMETRY

There are many parameters in our models. To reduce the parameters, we can impose a mirror [51] symmetry under which the fields transform as

$$\begin{aligned}&q'_L \leftrightarrow q_L\,,\ d'_R \leftrightarrow d_R\,,\ u'_R \leftrightarrow u_R\,,\ l'_L \leftrightarrow l_L\,,\ e'_R \leftrightarrow e_R\,,\\ &\phi' \leftrightarrow \phi\,,\ G'^a_\mu \leftrightarrow G^a_\mu\,,\ W'^a_\mu \leftrightarrow W^a_\mu\,,\ B'_\mu \leftrightarrow B_\mu\,,\\ &N'_R \leftrightarrow N_R\,,\ \Sigma_a \leftrightarrow \Sigma^T_a\,.\end{aligned}\tag{61}$$

The above mirror symmetry, which is assumed to softly break in the scalar potential, i.e.

$$\mu^2_{\phi'} \neq \mu^2_\phi\,,\ \mu^2_{\delta'} \neq \mu^2_\delta\,,\tag{62}$$

will simplify the parameters to be

$$\begin{aligned}&g'_3 = g_3\,,\ g'_2 = g_2\,,\ g'_1 = g_1\,,\ \lambda_{\phi'} = \lambda_\phi\,,\ \lambda_{\delta'} = \lambda_\delta\,,\\ &\lambda_{\phi\delta} = \lambda_{\phi'\delta'}\,,\ \lambda_{\phi\delta'} = \lambda_{\delta\phi'}\,,\ \lambda_{\phi\Sigma} = \lambda_{\phi'\Sigma}\,,\ y_{d'} = y_d\,,\\ &y_{u'} = y_u\,,\ y_{e'} = y_e\,,\ y_{\delta'} = y_\delta\,,\ y_{N'} = y_N\,,\ f_a = f^T_a\,,\\ &M_N = M^T_N\,.\end{aligned}\tag{63}$$

By fixing the VEVs:

$$\langle\phi\rangle' = 500\,\langle\phi\rangle\,,\tag{64}$$

we can read the dark charged fermion masses [50]:

$$\begin{array}{rclcrcl} m_{d'} &=& 2.5\,\mathrm{GeV} & \text{for} & m_d &=& 5\,\mathrm{MeV}\,,\\ m_{u'} &=& 1.25\,\mathrm{GeV} & \text{for} & m_u &=& 2.5\,\mathrm{MeV}\,,\\ m_{s'} &=& 50\,\mathrm{GeV} & \text{for} & m_s &=& 100\,\mathrm{MeV}\,,\\ m_{c'} &=& 645\,\mathrm{GeV} & \text{for} & m_c &=& 1.29\,\mathrm{GeV}\,,\\ m_{b'} &=& 2.095\,\mathrm{TeV} & \text{for} & m_b &=& 4.19\,\mathrm{GeV}\,,\\ m_{t'} &=& 86.45\,\mathrm{TeV} & \text{for} & m_t &=& 172.9\,\mathrm{GeV}\,,\\ m_{e'} &=& 0.256\,\mathrm{GeV} & \text{for} & m_e &=& 0.511\,\mathrm{MeV}\,,\\ m_{\mu'} &=& 52.85\,\mathrm{GeV} & \text{for} & m_\mu &=& 105.7\,\mathrm{MeV}\,,\\ m_{\tau'} &=& 888.5\,\mathrm{GeV} & \text{for} & m_\tau &=& 1.777\,\mathrm{GeV}\,.\end{array}\tag{65}$$

The beta functions of the QCD in the ordinary and dark sectors then will yield

$$\begin{aligned}\Lambda_{\mathrm{QCD}'} &= \left(\frac{\langle\phi'\rangle}{\langle\phi\rangle}\right)^{\frac{4}{11}} (m_u m_d m_s)^{\frac{2}{33}} \Lambda^{\frac{9}{11}}_{\mathrm{QCD}} = 1.13\,\mathrm{GeV}\\ &\quad \text{for}\ \ \Lambda_{\mathrm{QCD}} = 200\,\mathrm{MeV}\,.\end{aligned}\tag{66}$$

The dark proton and neutron masses thus can be given by

$$m_{p'} = 5\,\mathrm{GeV}\,,\ m_{n'} = 6.25\,\mathrm{GeV}\,.\tag{67}$$

In this case, the dark proton will serve as the dark matter particle.

Under the mirror symmetry, the Dirac neutrino mass matrices (22), (36) and (47) should be symmetric. Compared with the traditional type-I, type-II or type-I+II Majorana seesaw, our Dirac seesaw will not contain new parameters since the VEV in the dark sector has been determined by the dark matter mass.

VIII. SUMMARY

In this paper we have proposed a unified picture of the Dirac neutrino masses, the baryon asymmetry and the dark matter relic density. Specifically, we consider an $SU(3)'_c \times SU(2)'_L \times U(1)'_Y$ dark sector parallel to the $SU(3)_c \times SU(2)_L \times U(1)_Y$ ordinary sector and then introduce three types of messenger sectors composed of the heavy gauge-singlet Dirac fermion(s) and/or $[SU(2)_L \times SU(2)'_L]$-bidoublet Higgs scalar(s). Through the type-I, type-II or type-I+II Dirac seesaw mechanism, the heavy fermion singlet(s) and/or Higgs bidoublet(s) can highly suppress the masses between the ordinary and dark left-handed neutrinos. So, the ordinary and dark neutrinos can form the light Dirac neutrinos in a natural way. In such Dirac seesaw context, the lepton number conserving decays of the heavy fermion singlet(s) and/or Higgs bidoublet(s) can simultaneously generate a lepton asymmetry in the $[SU(2)_L]$-doublet leptons and an opposite lepton asymmetry in the $[SU(2)'_L]$-doublet leptons. Benefited from the $SU(2)_L$ and $SU(2)'_L$ sphaleron processes, we eventually can obtain a baryon asymmetry in the ordinary sector and an opposite baryon asymmetry in the dark sector. The lightest dark nucleon thus should have a determined mass about 5 GeV to serve as the dark matter particle. In the presence of the kinetic mixing between the $U(1)_Y$ and $U(1)'_Y$ gauge fields, the dark nucleons can be verified in the dark matter direct detection experiments. By imposing a mirror symmetry, our models needn't more parameters than the conventional Majorana seesaw models.

Acknowledgement: I thank Shao-Long Chen, Martin Holthausen, Manfred Lindner and Daniel Schmidt for discussions. This work is supported by the Sonderforschungsbereich TR 27 of the Deutsche Forschungsgemeinschaft.

---

[1] P. Minkowski, Phys. Lett. B **67**, 421 (1977); T. Yanagida, in *Proceedings of the Workshop on Unified Theory and the Baryon Number of the Universe*, edited by O. Sawada and A. Sugamoto (KEK, Tsukuba, 1979), p. 95; M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by F. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979), p. 315; S.L. Glashow, in *Quarks and Leptons*, edited by M. Lévy *et al.* (Plenum, New York, 1980), p. 707; R.N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).

[2] M. Magg and C. Wetterich, Phys. Lett. B **94**, 61 (1980); J. Schechter and J.W.F. Valle, Phys. Rev. D **22**, 2227 (1980); T.P. Cheng and L.F. Li, Phys. Rev. D **22**, 2860 (1980); G. Lazarides, Q. Shafi, and C. Wetterich, Nucl. Phys. B **181**, 287 (1981); R.N. Mohapatra and G. Senjanović, Phys. Rev. D **23**, 165 (1981).

[3] R. Foot, H. Lew, X.G. He, and G.C. Joshi, Z. Phys. C **44**, 441 (1989).

[4] M. Roncadelli and D. Wyler, Phys. Lett. B **133**, 325 (1983); P. Roy and O. Shanker, Phys. Rev. Lett. **52**, 713 (1984).

[5] H. Murayama and A. Pierce, Phys. Rev. Lett. **89**, 271601 (2002).

[6] P.H. Gu and H.J. He, JCAP **0612**, 010 (2006).

[7] M. Fukugita and T. Yanagida, Phys. Lett. B **174**, 45 (1986).

[8] R.N. Mohapatra and X. Zhang, Phys. Rev. D **46**, 5331 (1992); E. Ma and U. Sarkar, Phys. Rev. Lett. **80**, 5716 (1998).

[9] M.Flanz, E.A. Paschos, and U. Sarkar, Phys. Lett. B **345**, 248 (1995); M.Flanz, E.A. Paschos, U. Sarkar, and J. Weiss, Phys. Lett. B **389**, 693 (1996); L. Covi, E. Roulet, and F. Vissani, Phys. Lett. B **384**, 169 (1996); A. Pilaftsis, Phys. Rev. D **56**, 5431 (1997).

[10] S. Davidson and A. Ibarra, Phys. Lett. B **535**, 25 (2002); W. Buchmüller, P. Di Bari, and M. Plümacher, Nucl. Phys. B **665**, 445 (2003).

[11] T. Hambye and G. Senjanović, Phys. Lett. B **582**, 73 (2004); S. Antusch and S.F. King, Phys. Lett. B **597**, 199 (2004).

[12] T. Hambye, Y. Lin, A. Notari, M. Papucci, A. Strumia, Nucl. Phys. B **695**, 169 (2004).

[13] S. Blanchet and P. Di Bari, Nucl. Phys. B **807**, 155 (2009).

[14] P.H. Gu, Phys. Lett. B **713**, 485 (2012).

[15] K. Dick, M. Lindner, M. Ratz, and D. Wright, Phys. Rev. Lett. **84**, 4039 (2000).

[16] B. Thomas and M. Toharia, Phys. Rev. D **73**, 063512 (2006); S. Abel and V. Page, JHEP **0605**, 024 (2006); D.G. Cerdeno, A. Dedes, and T.E.J. Underwood, JHEP **0609**, 067 (2006); E.J. Chun and P. Roy, JHEP **0806**, 089 (2008); P.H. Gu, H.J. He, and U. Sarkar, Phys. Lett. B **659**, 634 (2008); A. Bechinger and G. Seidl, Phys. Rev. D **81**, 065015 (2010); H. Davoudiasl and I. Lewis, arXiv:1112.1939 [hep-ph].

[17] E. Komatsu *et al.*, Astrophys. J. Suppl. **192**, 18 (2011).

[18] S. Nussinov, Phys. Lett. B **165**, 55 (1985).

[19] S.M. Barr, R.S. Chivulula, and E. Farhi, Phys. Lett. B **241**, 387 (1990); S.M. Barr, Phys. Rev. D **44**, 3062 (1991).

[20] D.B. Kaplan, Phys. Rev. Lett. **68**, 741 (1992).

[21] S. Dodelson, B.R. Greene, and L.M. Widrow, Nucl. Phys. B **372**, 467 (1992).

[22] V.A. Kuzmin, Phys. Part. Nucl. **29**, 257 (1998) [Fiz. Elem. Chast. Atom. Yadra **29**, 637 (1998)].

[23] R. Kitano and I. Low, Phys. Rev. D **71**, 023510 (2005).

[24] K. Agashe and G. Servant, JCAP **0502**, 002 (2005); M. Cirelli, P. Panci, G. Servant, and G. Zaharijas, JCAP **1203**, 015 (2012).

[25] N. Cosme, L. Lopez Honorez, and M.H.G. Tytgat, Phys. Rev. D **72**, 043505 (2005).

[26] P.H. Gu, U. Sarkar, and X. Zhang, Phys. Rev. D **80**, 076003 (2009); P.H. Gu and U. Sarkar, Phys. Rev. D **81**, 033001 (2010); P.H. Gu, M. Lindner, U. Sarkar, and X. Zhang, Phys. Rev. D **83**, 055008 (2011).

[27] H. An, S.L. Chen, R.N. Mohapatra, and Y. Zhang, JHEP **1003**, 124 (2010); H. An, S.L. Chen, R.N. Mohapatra, S. Nussinov, and Y. Zhang, Phys. Rev. D **82**, 023533 (2010).

[28] H. Davoudiasl, D.E. Morrissey, K. Sigurdson, and S. Tulin, Phys. Rev. Lett. **105**, 211304 (2010); H. Davoudiasl, D.E. Morrissey, K. Sigurdson, and S. Tulin, Phys. Rev. D **84**, 096008 (2011); N. Blinov, D.E. Morrissey, and S. Tulin, arXiv:1206.3304 [hep-ph].

[29] M. Blennow, B. Dasgupta, E. Fernandez-Martinez, and N. Rius, JHEP **1103**, 014 (2011); M. Blennow, E. Fernandez-Martinez, J. Rendondo, and P. Serra, arXiv:1203.5805 [hep-ph].

[30] J. McDonald, Phys. Rev. D **83**, 083509 (2011); Phys. Rev. D **84**, 103514 (2011).

[31] L.J. Hall, J. March-Russell, and S.M. West, arXiv:1010.0245 [hep-ph]; J. March-Russell, M. McCullough, JCAP **1203**, 019 (2012); J. March-Russell, J. Unwin, and S.M. West, arXiv:1203.4854 [hep-ph].

[32] B. Dutta and J. Kumar, Phys. Lett. B **699**, 364 (2011).

[33] A. Falkowski, J.T. Ruderman, and T. Volansky, JHEP **1105**, 106 (2011).

[34] N. Haba, S. Matsumoto, and R. Sato, Phys. Rev. D **84**, 055016 (2011).

[35] Z. Kang, J. Li, T. Li, T. Liu, and J. Yang, arXiv:1102.5644 [hep-ph]; Z. Kang and T. Li, arXiv:1111.7313 [hep-ph].

[36] M.L. Graesser, I.M. Shoemaker, and L. Vecchi, JHEP **1110**, 110 (2011).

[37] M.T. Frandsen, S. Sarkar, and K. Schmidt-Hoberg, Phys. Rev. D **84**, 051703 (2011).

[38] S.D. McDermott, H.B. Yu, and K.M. Zurek, Phys. Rev. D **85**, 023519 (2012); S. Tulin, H.B. Yu, and K.M. Zurek, JCAP **1205**, 013 (2012).

[39] H. Iminniyaz, M. Drees, and X. Chen, JCAP **107**, 003 (2011).

[40] N.F. Bell, K. Petraki, I.M. Shoemaker, and R.R. Volkas, Phys. Rev. D **84**, 123505 (2011); K. Petraki, M. Trodden, and R.R. Volkas, JCAP **1202**, 044 (2012); B. von Harling, K. Petraki, and R.R. Volkas, JCAP **1205**, 021 (2012).

[41] Y. Cui, L. Randall, and B. Shuve, JHEP **1108**, 073 (2011); Y. Cui, L. Randall, and B. Shuve, JHEP **1204**, 075 (2012).

[42] C. Arina, and N. Sahu, Nucl. Phys. B **854**, 666 (2012); C. Arina, J.O. Gong, and N. Sahu, arXiv: 1206.0009 [hep-ph].

[43] E. Ma and U. Sarkar, arXiv:1111.5350 [hep-ph].

[44] H. Davoudiasl and R.N. Mohapatra, arXiv:1203.1247 [hep-ph].

[45] W.Z. Feng, P. Nath, and G. Peim, arXiv:1204.5752 [hep-ph].

[46] G. t'Hooft, Phys. Rev. Lett. **37**, 8 (1976); Phys. Rev. D **14**, 3432 (1976).

[47] V.A. Kuzmin, V.A. Rubakov, and M.E. Shaposhnikov, Phys. Lett. B **155**, 36 (1985).

[48] T. Asaka, S. Blanchet, and M. Shaposhnikov, Phys. Lett. B **631**, 151 (2005); A. Boyarsky, O. Ruchayskiy, and M. Shaposhnikov, Ann. Rev. Nucl. Part. Sci. **59**, 191 (2009).

[49] E.W. Kolb and M.S. Turner, *The Early Universe*, Addison-Wesley, 1990.

[50] K. Nakamura *et al.*, (Particle Data Group), J. Phys. G **37**, 075021 (2010).

[51] T.D. Lee and C.N. Yang, Phys. Rev. textbf104, 254 (1956); I.Yu. Kobzarev, L.B. Okun, and I.Ya. Pomeranchuk, Sov. J. Nucl. Phys. **3**, 837 (1966) [Yad. Fiz. **3**, 1154 (1966)]; M. Pavsic, Int. J. Theor. Phys. **9**, 229 (1974); S.I. Blinnikov and M.Y. Khlopov, Sov. J. Nucl. Phys. **36**, 472 (1982) [Yad. Fiz. **36**, 809 (1982)]; S. L. Glashow, Phys. Lett. B **167**, 35 (1986); R. Foot, H. Lew, and R.R. Volkas, Phys. Lett. B **272**, 67 (1991); H.M. Hodges, Phys. Rev. D **47** 456 (1993); R. Foot and R.R Volkas, Phys. Rev. D **52**, 6595 (1995); Z. Berezhiani and R.N. Mohapatra, Phys. Rev. D **52**, 6607 (1995); Z. Berezhiani, A. Dolgov, and R.N. Mohapatra, Phys. Lett. B **375**, 26 (1996); R.N. Mohapatra and V.L. Teplitz, Phys. Lett. B **462**, 302 (1999); Phys. Rev. D **62**, 063506 (2000); Z. Berezhiani, D. Comelli, and F.L. Villante, Phys. Lett. B **503** 362 (2001); R.N. Mohapatra, S. Nussinov and V.L. Teplitz, Phys. Rev. D **66**, 063002 (2002); A.Y. Ignatiev and R.R. Volkas, Phys. Rev. D **68**, 023518 (2003); Z. Berezhiani, P. Ciarcelluti, D. Comelli, and F. L. Villante, Int. J. Mod. Phys. D 14, 107 (2005); P. Ciarcelluti, Int. J. Mod. Phys. D 14, 187 (2005); C.R. Das, L.V. Laperashvili, H.B. Nielsen, and A. Tureanu, Phys. Rev. D **84**, 063510 (2011); J.W. Cui, H.J. He, L.C. Lü, and F.R. Yin, Phys. Rev. D **85**, 096003 (2012).